



FastFung: A novel medium for the culture and isolation of fastidious fungal species from clinical samples

Fadi Bittar^{a,c}, Frédérique Gouriet^{a,c}, Saber Khelaifia^{a,c}, Didier Raoult^{a,c}, Stéphane Ranque^{b,c,*}

^a Aix Marseille Univ, IRD, APHM, MEPHI, Marseille, France

^b Aix Marseille Univ, IRD, APHM, SSA, VITROME, Marseille, France

^c IHU Méditerranée Infection, Marseille, France

ARTICLE INFO

Keywords:

Culture media
Clinical laboratory techniques
Mycology
Mycoses
Malassezia
Candida
Cryptococcus
Mycobiome

ABSTRACT

We developed a novel culture medium, referred to FastFung medium as suitable for the culture of clinical fungi, including fastidious ones, for both research and diagnostic studies. It is based on Schaedler agar supplemented with many essential components for the growth of fastidious fungi. It also contains selective antibacterial agents for the inhibition of contaminant bacteria growth. In this preliminary study, the FastFung medium was compared to the gold standard Sabouraud medium for 98 fungal and 20 bacterial strains.

The fungal strain positive culture rate was 100% vs. 95% and the bacterial strain inhibition was 100% vs. 20%, for the FastFung and Sabouraud media, respectively. When compared to the Sabouraud medium on 120 clinical samples, the FastFung medium displayed both a higher fungal colonies count, and a lower culture contamination rate. Storage at 4 °C for 4 weeks did not alter the FastFung culture medium performances for the six isolates of *Candida*, *Cryptococcus*, and *Penicillium* tested.

These encouraging results suggest future development of using the FastFung medium in clinical mycology and in mycobiome characterization. Further prospective evaluation aiming at assessing whether implementing the FastFung medium in the routine workflow simplifies and strengthens fungal isolation capacities in the clinical laboratory is warranted.

1. Introduction

Fungal species are ubiquitous environmental organisms. They can be found in food, water, soil and also isolated from humans and animals samples (Praneenararat, 2014). They are often associated with human infectious diseases (Botnaru et al., 2014; Rodriguez and Ramos, 2014). Fungal microorganisms of clinical interest are mainly grouped into three classes: Ascomycota, Basidiomycota and Zygomycota (Gouba & Drancourt, 2015). Fungal infections are the current object of many studies (Praneenararat, 2014; Botnaru et al., 2014; Chen et al., 2011). The predominant clinical species is *Candida albicans* that is involved in the majority of opportunistic invasive fungal diseases (Vidya et al., 2016). Indeed, recent studies have pointed out *C. albicans* as a major cause of morbidity and mortality in critical care patients, as well as its involvement in other oropharyngeal diseases of the mouth, and digestive and vaginal tract in immunocompromised and antibiotic treated individuals (Huffnagle & Noverr, 2013). The genera *Cryptococcus* and *Aspergillus* also include many fungi involved in human infections (Botnaru et al.,

2014; Geltner & Lass-Flörl, 2016). Many fungal species are opportunistic pathogens; i.e. they are commensals when the host is healthy, and they become pathogenic when the host is immunocompromised and/or critically ill (Huffnagle & Noverr, 2013). Thus, the population of immunocompromised patients has the highest risk of opportunistic fungal infections (Botnaru et al., 2014; Rodriguez & Ramos, 2014; de Oliveira et al., 2014). Moreover, recent increases in emerging fungal infections call for an effective culture medium capable of timely isolating fastidious fungi from any biological sample at the clinical mycology laboratory.

While Sabouraud dextrose agar is the most frequently used medium for fungal isolation (Gumral et al., 2015; Matic Petrović et al., 2015), other media are commonly used for culture of yeasts and filamentous fungi, including Potatoe Dextrose Agar (Gouba et al., 2013), Czapek (Ali et al., 2014) and Dixon agar (Shah et al., 2013). These media are often used in combination in clinical laboratories. These, and several other culture media, are used as reference for the isolation or the identification of specific fungal taxa (Ben Salah et al., 2010). This study aimed at

* Corresponding author at: Parasitologie – Mycologie, IHU Méditerranée Infection, 19-21 Boulevard Jean Moulin, 13005 Marseille, France.

E-mail address: stephane.ranque@ap-hm.fr (S. Ranque).

<https://doi.org/10.1016/j.mimet.2020.106108>

Received 6 August 2020; Received in revised form 17 November 2020; Accepted 18 November 2020

Available online 21 November 2020

0167-7012/© 2020 Elsevier B.V. All rights reserved.

Table 1

Culture success rates of 98 pure culture clinical fungal isolates inoculated onto Sabouraud and FastFung media.

Fungal species	N	Sabouraud	FastFung
<i>Acremonium kiliense</i>	2	100%	100%
<i>Aspergillus flavus</i>	1	100%	100%
<i>Aspergillus niger</i>	5	100%	100%
<i>Aspergillus sydowii</i>	1	100%	100%
<i>Beauveria bassiana</i>	2	100%	100%
<i>Candida albicans</i>	10	100%	100%
<i>Candida glabrata</i>	6	100%	100%
<i>Candida guilliermondii</i>	5	100%	100%
<i>Candida kefyr</i>	4	100%	100%
<i>Candida lusitanae</i>	3	100%	100%
<i>Candida orthopsilosis</i>	6	100%	100%
<i>Candida parapsilosis</i>	6	100%	100%
<i>Candida tropicalis</i>	6	100%	100%
<i>Candida zeylanoides</i>	1	0%	100%
<i>Candida krusei</i>	3	100%	100%
<i>Clavispora lusitanae</i>	2	100%	100%
<i>Cryptococcus diffluens</i>	1	0%	100%
<i>Cryptococcus unigutulatus</i>	1	0%	100%
<i>Debaryomyces hansenii</i>	2	100%	100%
<i>Galactomyces geotrichum</i>	1	100%	100%
<i>Isaria farinosa</i>	1	100%	100%
<i>Malassezia furfur</i>	1	0%	100%
<i>Malassezia pachydermatis</i>	1	100%	100%
<i>Mucor circinelloides</i>	1	100%	100%
<i>Mucor velutinosus</i>	1	100%	100%
<i>Penicillium chrysogenum</i>	1	100%	100%
<i>Penicillium glandicola</i>	1	100%	100%
<i>Penicillium dipodomyicola</i>	1	100%	100%
<i>Pichia caribbica</i>	3	100%	100%
<i>Pichia manshurica</i>	1	100%	100%
<i>Rhodotorula mucilaginosa</i>	6	100%	100%
<i>Saccharomyces cerevisiae</i>	6	100%	100%
<i>Trichosporon asahii</i>	4	100%	100%
<i>Preussia minima</i>	1	100%	100%
<i>Torulaspota pretoriensis</i>	1	0%	100%
Total	98	95%	100%

developing a novel culture medium, which is suitable for the culture of clinical fungi, including fastidious ones, for both research and diagnostic studies. The FastFung medium is based on Schädler agar (Sigma-Aldrich) supplemented with many components needed for the growth of fungal species. It also contains selective antibacterial agents for the inhibition of contaminant bacteria. In this preliminary study, the FastFung medium was evaluated on 98 fungal and 20 bacterial strains.

2. Materials and methods

2.1. Culture media

In this study the performance of fungal culture on FastFung and Sabouraud dextrose agar (Oxoid, Dardilly, France) media were compared. The Sabouraud medium is supplemented with chloramphenicol and gentamycin. In addition to bacteriological agar, this culture medium contains glucose and peptone allowing the growth of many fungal species. Sabouraud medium was considered as the gold-standard fungal medium in this study. The FastFung medium proposed in this study is composed per liter of 43 g of Schädler Agar (Sigma-Aldrich, Saint-Quentin Fallavier, France), 20 g of peptone and 10 g of glucose as basic components, with the addition of 5 g of ox-bile 10 g of malt extract, 2 ml of oleic acid, 5 ml of Tween 60 and 2.5 ml of glycerol. Each component was autoclaved at 121 °C for 30 min; except for Tween 60, glycerol and oleic acid, which were added after autoclaving. After cooling to 50 °C, the FastFung medium was supplemented with a mixture of three antibiotics at 30 mg/l, including colistin, vancomycin and imipenem (Sigma Aldrich, St Louis, France). The final product was then dispensed into 90 mm sterile Petri dishes.

Table 2

Isolation rates of pure bacterial strains inoculated onto Sabouraud and FastFung media.

Bacterial strains	N	Sabouraud	FastFung
<i>Acinetobacter baumannii</i>	2	100%	0%
<i>Enterobacter cloacae</i>	2	100%	0%
<i>Enterococcus faecalis</i>	2	100%	0%
<i>Escherichia coli</i>	2	100%	0%
<i>Gardnerella vaginalis</i>	2	100%	0%
<i>Haemophilus influenzae</i>	2	100%	0%
<i>Klebsiella oxytoca</i>	2	100%	0%
<i>Klebsiella pneumoniae</i>	2	100%	0%
<i>Proteus mirabilis</i>	1	0%	0%
<i>Pseudomonas aeruginosa</i>	1	0%	0%
<i>Rhizobium radiobacter</i>	1	0%	0%
<i>Staphylococcus aureus</i>	1	0%	0%
Total	20	80%	0%

Table 3

Fungi cultivation rates from various routine clinical samples inoculated onto Sabouraud and FastFung media.

Samples	Species	N	Sabouraud	FastFung
Vaginal	<i>Candida albicans</i>	52	100%	100%
Urine	<i>Candida albicans</i>	70	100%	100%
Urine	<i>Candida tropicalis</i>	15	100%	100%
Swab	<i>Candida albicans</i>	4	100%	100%
Nasopharyngeal aspiration	<i>Candida albicans</i>	4	100%	100%
Bronchial aspiration	<i>Candida tropicalis</i>	2	100%	100%
Bronchial aspiration	<i>Candida parapsilosis</i>	3	100%	100%
Sputum	<i>Candida albicans</i>	6	100%	100%
Pulmonary aspiration	<i>Candida albicans</i>	2	100%	100%
Aortic valve	<i>Candida albicans</i>	2	100%	100%
Biopsy	<i>Candida albicans</i>	1	100%	100%
Total		161	100%	100%

2.2. Bacterial and fungal strains

A collection of 98 fungal strains (Table 1) and 20 bacterial strains (Table 2) that were isolated in the laboratory by culturomics methods (Lagier et al., 2012, 2015) were used to assess the FastFung medium. The identification of each strain was confirmed by the sequence analysis of the partial 16S rRNA gene for bacteria or the ITS rRNA region for fungi, as previously described (Lagier et al., 2012; Hamad et al., 2012). The fungal strains used included *Candida* spp. ($n = 50$), *Cryptococcus* spp. ($n = 2$), *Aspergillus* spp. ($n = 7$), *Penicillium* spp. ($n = 3$), *Malassezia* spp. ($n = 2$) and 34 other fungal strains (Table 1). Appropriate negative and positive controls were used in each experiment.

2.3. Evaluation of FastFung medium and Sabouraud agar on fungal strains

All the strains used in this study have been preserved at -80 °C in sterile tubes with glycerol. The fungal strains have been cultured onto Sabouraud, Dixon agar or Potatoes Dextrose agar and the bacterial strains on 5% sheep blood-enriched Colombia agar (BioMérieux, France) to obtain fresh and pure cultures. Each isolate was suspended in 1 ml of saline (0.85%) at a concentration of approximately 1.5×10^5 CFU/ml for fungi and 1.5×10^8 CFU/ml for bacteria (0.5 McFarland standard) using a densitometer (DEN-1 McFarland Densitometer, BioSan, Riga, Latvia). Serial five-fold inoculum dilutions were then prepared as follows: 100 μ l of each solution was diluted in 900 μ l of water following by four other serial dilutions. A 1 μ l aliquot of each dilution was inoculated onto the culture plate and spread with a sterile inoculation loop. All plates were incubated at 28 °C and 32 °C for fungi and 37 °C for bacterial strains under aerobic conditions. The bacterial strains were inoculated in the same way as fungal strains, but no dilution was done.




Fig. 1. The FastFung medium displayed higher fungal CFU counts compared to the Sabouraud medium. Here we show the isolation after 48 h of incubation of three yeast species (*Candida albicans*, *C. parapsilosis*, and *C. tropicalis*) from routine clinical samples that were inoculated in parallel onto both culture media.

2.4. Assessment of the FastFung medium on routine clinical samples

The effectiveness of the FastFung medium was further assessed on 120 clinical samples, prospectively received at the Timone University Hospital (Marseille, France) microbiology laboratory for routine analyses for a 6 week period in which a fungus was isolated. These samples were obtained from various body sites, mostly urine and vagina, as detailed in Table 3. An aliquot (50 μ l) of each clinical sample was inoculated onto the Sabouraud and the FastFung media in parallel. The plates were incubated under aerobic conditions from 24 to 48 h at 28 °C.

2.5. FastFung medium stability assessment

Forty FastFung medium plates were prepared and stored at 4 °C for 4 weeks. Six fungal strains including *Candida* spp. ($n = 2$), *Cryptococcus* spp. ($n = 2$) and *Penicillium* spp. ($n = 2$), were used to test the stability of the culture medium. The same procedures described above were applied to inoculate fungal strains onto the FastFung medium. Inoculated plates were then incubated under aerobic conditions at 28 °C and examined daily for 7 days.

3. Results

3.1. FastFung medium assessment on fungal strains

The positive culture rate of the fungal strains after 6 days 98/98 (100%) with the FastFung medium. Moreover, each inoculum dilution yielded a positive culture with the FastFung medium. Among each fungal strain assayed on the FastFung medium, growth was visualized in 57/98 (58%) of the cultures after 24 h of incubation and in 67/98 (68%) after 48 h of incubation. In contrast, the Sabouraud medium allowed the growth of 93/98 (95%) of the fungal strains, and in particular, *Candida zeylanoides*, *Cryptococcus diffluens*, *Cryptococcus unigutulatus*, *Malassezia furfur* and *Torulasporea pretoriensis* did not grow (Table 1). Among bacterial strains inoculated onto the FastFung medium plates, 20/20

(100%) bacterial strains were effectively inhibited and did not grow on the FastFung medium (Table 2). In contrast, only 4/20 (20%) of the bacterial strains were inhibited and did not grow on the Sabouraud medium (Table 2).

3.2. FastFung medium assessment on prospective routine clinical samples

There was no discrepancy between the two media regarding the positive culture rate. All the tested were culture positive samples grew on both Sabouraud and FastFung culture media (Table 3). *Candida albicans*, *Candida tropicalis* and *Candida parapsilosis* were the most frequently isolated fungal species. *Candida albicans* was the most frequent isolated species, regardless of the clinical sample type and the culture medium. *Candida albicans* and *Candida tropicalis* were relatively more frequently isolated from urinary samples. *Candida tropicalis* and *Candida parapsilosis* were relatively more frequently isolated from bronchial aspiration samples. After only 24 h of incubation the number of fungal CFU was higher on the FastFung medium than on the Sabouraud medium (Fig. 1). Moreover, several filamentous fungi contaminations occurred on Sabouraud but not on FastFung medium (data not shown).

3.3. FastFung medium stability

Neither the quality of growth of the six fungal strains nor the inhibition bacterial strains and filamentous fungi was altered when the culture medium had been stored at 4 °C for 4 weeks (data not shown).

4. Discussion

Overall the FastFung medium proved more efficient than the Sabouraud medium for the rapid isolation of fungi and inhibition of bacteria and filamentous fungi contaminants from the clinical samples tested. Each of the 98 fungal strains tested grew on the FastFung medium, whereas only 67 grew on Sabouraud. A number of fastidious fungi, such

as the lipid dependent *Malassezia* spp. yeast, require the specific culture media for their growth (Shah et al. 2013, Ben Salah et al. 2010). The addition of specific components, such as oleic acid, considerably extended the spectra of culturable fungi as described previously (Mayser et al., 1997). In addition, and although both two culture media contain antibacterials, the FastFung medium more efficiently inhibited bacterial growth than the Sabouraud medium.

Studies comparing the efficiency of different fungal culture media are scarce. To our knowledge, we report the first study that compares a new fungal culture medium with a reference culture medium commonly used for the isolation and culture of fungal species. One strengths of this study is the large number of bacterial and fungal strains assessed, which were all identified by MALDI TOF mass spectrometry and DNA sequence analysis (Sow et al., 2015).

When compared to the Sabouraud medium on routine clinical samples, FastFung medium displayed a higher fungal colonies count after 48 h of culture (Fig. 1) a lower number of commonly encountered culture contamination (Gumral et al., 2015; Sharma et al., 2014). The FastFung medium proved stable for 4 weeks. Using this versatile culture medium in the clinical laboratory would advantageously avoid using various fungal culture media for the isolation of fastidious fungi and offers great opportunities in clinical microbiology and culturomics studies when associated with MALDI TOF mass spectrometry identification based on comprehensive fungal reference spectra databases.

In conclusion, implementing the FastFung medium in the routine clinical mycology laboratory workflow would advantageously reduce the time to fungal growth and makes unnecessary the use of Sabouraud and other complementary culture media for fungal isolation. Further prospective evaluation of the use of the FastFung medium for fungal isolation in the clinical laboratory is warranted.

Ethical approval

Not required.

Funding information

This work was supported by the French Government under the «Investissements d'avenir» (Investments for the Future) program managed by the Agence Nationale de la Recherche (ANR, fr: National Agency for Research), (reference: Méditerranée Infection 10-IAHU-03).

This work was supported by Région Provence-Alpes-Côte d'Azur and European funding FEDER (fonds européen de développement régional) PRIMMI ((Plateformes de Recherche et d'Innovation Mutualisées Méditerranée Infection).

Declaration of Competing Interest

None of the authors have competing interest to declare.

Acknowledgements

This work is dedicated to the memory of Marcel Sy (PhD student died

on 15 September 2016); Marcel Sy has made a substantial contribution to this work.

References

- Ali, S., Malik, A., Bhargava, R., Shahid, M., Fatima, N., 2014. Aspergillus colonization in patients with bronchogenic carcinoma. *Asian Cardiovasc. Thorac. Ann.* 22 (4), 460–464.
- Ben Salah, I., Makni, F., Cheikhrouhou, F., Neji, S., Sellami, H., Ayadi, A., 2010. *Malassezia* species: pathology, isolation and identification media. *J. Mycol. Med.* 20, 53–60. <https://doi.org/10.1016/j.mycmed.2009.11.006>.
- Botnaru, V., Rusu, D., Haidarli, I., Munteanu, O., Corlateanu, A., 2014. Cryptococcosis–infectie fungica frecventa a imunocompromisului Cryptococcosis-a common fungal infection in immunosuppressed patient. *Pneumologia* 63 (3), 159–163, 156. Romanian. PMID: 25420290.
- Chen, Y., Chen, Z., Guo, R., Chen, N., Lu, H., Huang, S., et al., 2011. Correlation between gastrointestinal fungi and varying degrees of chronic hepatitis B virus infection. *Diagn. Microbiol. Infect. Dis.* 70 (4), 492–498.
- Geltner, C., Lass-Flörl, C., 2016. Invasive pulmonary Aspergillosis in organ transplants - focus on lung transplants. *Respir. Investig.* 54 (2), 76–84.
- Gouba, N., Drancourt, M., 2015. Digestive tract mycobiota: a source of infection. *Med. Mal. Infect.* 45 (1–2), 9–16.
- Gouba, N., Raoult, D., Drancourt, M., 2013. Plant and fungal diversity in gut microbiota as revealed by molecular and culture investigations. *PLoS One* 8 (3), e59474.
- Gumral, R., Dogen, A., Ilkit, M.M., 2015. Comparison of the contamination rates of culture media used for isolation and identification of dermatophytes. *Turk J Med Sci* 45 (3), 587–592.
- Hamad, I., Sokhna, C., Raoult, D., Bittar, F., 2012. Molecular detection of eukaryotes in a single human stool sample from Senegal. *PLoS One* 7 (7), e40888.
- Huffnagle, G.B., Noverr, M.C., 2013. The emerging world of the fungal microbiome. *Trends Microbiol.* 21 (7), 334–341.
- Lagier, J.C., Armougou, F., Million, M., Hugon, P., Pagnier, I., Robert, C., et al., 2012. Microbial culturomics: paradigm shift in the human gut microbiome study. *Clin. Microbiol. Infect.* 18 (12), 1185–1193.
- Lagier, J.C., Hugon, P., Khelaifa, S., Fournier, P.E., La, S.B., Raoult, D., 2015. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin. Microbiol. Rev.* 28 (1), 237–264.
- Matić Petrović, S., Cimbajević, M., Radunović, M., Kuzmanović Pfićer, J., Jotić, A., Pucar, A., 2015. Detection and sampling methods for isolation of *Candida* spp. from oral cavities in diabetics and non-diabetics. *Braz. Oral Res.* 29 <https://doi.org/10.1590/1807-3107BOR-2015.S1806-83242015000100272.vol29.0077>. PMID: (26083092).
- Mayser, P., Haze, P., Papavassilis, C., Pickel, M., Gruender, K., Gueho, E., 1997. Differentiation of *Malassezia* species: selectivity of cremophor EL, castor oil and ricinoleic acid for *M. furfur*. *Br. J. Dermatol.* 137 (2), 208–213.
- de Oliveira, R.B., Atobe, J.H., Souza, S.A., de Castro Lima Santos, D.W., 2014. Epidemiology of invasive fungal infections in patients with acquired immunodeficiency syndrome at a reference hospital for infectious diseases in Brazil. *Mycopathologia* 178 (1–2), 71–78.
- Praneenarat, S., 2014. Fungal infection of the colon. *Clin. Exp. Gastroenterol.* 7, 415–426.
- Rodriguez, Z.C., Ramos, M.G., 2014. Acremonium species associated fungemia: a novel pathogen in the immunosuppressed patient. *Bol. Asoc. Med. P. R.* 106 (3), 29–31.
- Shah, A., Koticha, A., Ubale, M., Wanjare, S., Mehta, P., Khopkar, U., 2013. Identification and speciation of malassezia in patients clinically suspected of having pityriasis versicolor. *Indian J. Dermatol.* 58 (3), 239.
- Sharma, P., Kumar, N., Jain, S., 2014. Fungal contaminants in cytopathology specimens. *Biom. J.* 37 (1), 31–32.
- Sow, D., Fall, B., Ndiaye, M., Ba, B.S., Sylla, K., Tine, R., et al., 2015. Usefulness of MALDI-TOF mass spectrometry for routine identification of *Candida* species in a resource-poor setting. *Mycopathologia* 180 (3–4), 173–179.
- Vidya, K.M., Rao, U.K., Nittayananta, W., Liu, H., Owotade, F.J., 2016. Oral mycoses and other opportunistic infections in HIV: therapy and emerging problems - a workshop report. *Oral Dis.* 22 (Suppl. 1), 158–165.